

BOEHRINGER INGELHEIM (CANADA) LTD.

5180 SOUTH SERVICE ROAD, BURLINGTON, ON, L7L 5H4

PRODUCT INFORMATION NO. 1-800-263-5103
TECHNICAL SERVICES NO. 1-877-565-5501
WEBSITE: www.boehringer-ingelheim.ca



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the Canada product label or package insert.

BOVIKALC®

BOEHRINGER

Veterinary Use Only

Oral calcium supplement for freshening cows

Warnings: No withdrawal period for meat and withholding time for milk are required when used according to the label. Keep out of reach of children.

NN.S0Z9

FAST & SUSTAINED RELEASE OF CALCIUM

Bovikalc® is an oral calcium supplement for use around parturition in fresh dairy cows.

Direction for Use: Minimum Protocol: Give one bolus immediately after calving and a second bolus 12 - 24 hours after calving. Only administer using the Bovikalc® branded applicator.

Extended Protocol: For high yielding cows administer:

- 1st bolus at the first signs of parturition
- 2nd bolus immediately after calving
- 3rd bolus 12 - 24 hours after calving
- 4th bolus 24 - 48 hours after calving

Please discuss protocol with your veterinarian before implementation.

Place the bolus in the bolus applicator with the rounded end pointing outward.

If right-handed, place yourself at the cow's right side.

Grab the cow's upper jaw with your left hand. Keep the cow's head at a natural position and a little towards yourself. Gently guide the loaded Bovikalc® applicator into the cow's mouth.

When you feel resistance, gently guide the applicator over the tongue. When the applicator is behind the tongue, squeeze the handle to release the bolus.

Wait a moment for the cow to swallow and then remove the applicator.

Active Ingredients per bolus: Calcium chloride - 579.1 mg/g, Calcium sulfate - 231.6 mg/g

Inactive Ingredients: Water, Acetylated monoglycerides, Xanthan gum

Contraindications: Not for use in cows for initial treatment of milk fever, with clinical signs of milk fever or laying cows.

Do not use broken or damaged boluses.

Store at or below 30°C. Boluses are sensitive to humidity and should be stored in the plastic tubes until immediately before use.

Bovikalc® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH, used under license.

ATTENTION! DO NOT USE BROKEN OR DAMAGED BOLUSES.



MANUFACTURED BY: Boehringer Ingelheim Denmark A/S, DK-2100 Copenhagen OE, 208-R854833

MADE IN DENMARK

NET CONTENTS: 4 x 192 g BOLUSES 125887-001

CPN: 1230152.0

Bovikalc® is a registered trademark of Boehringer Ingelheim Vetmedica GmbH, used under license.